



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa


Uniqueness of symmetric basis in quasi-Banach spaces

F. Albiac*, C. Leránoz

Departamento de Matemáticas, Universidad Pública de Navarra, Pamplona 31006, Spain

ARTICLE INFO

Article history:

Received 10 June 2008

Available online 11 July 2008

Submitted by J. Diestel

Keywords:

Quasi-Banach space

Symmetric basis

ABSTRACT

We show that if X is a nonlocally convex natural quasi-Banach space with symmetric basis whose Banach envelope is isomorphic to ℓ_1 , then all symmetric bases of X are equivalent. The scope of this result is quite ample since the Banach envelopes of natural quasi-Banach spaces with basis always exhibit an ℓ_1 -like behavior, in the sense that they contain copies of ℓ_1^n 's uniformly complemented.

© 2008 Elsevier Inc. All rights reserved.

1. Introduction

A quasi-Banach space X is said to have a unique unconditional or symmetric basis if for any two normalized bases of the same type, there exists an automorphism of X which takes one basis to the other.

If X is locally convex (i.e., X is Banach space) then X has a unique unconditional basis if and only if it is isomorphic to one of the spaces ℓ_2 , ℓ_1 or c_0 [17]. The class of Banach spaces with a unique symmetric basis is considerably larger and it contains all the Orlicz sequence spaces ℓ_F for which $\lim_{t \rightarrow 0} tF'(t)/F(t)$ exists [15], and also the Lorentz sequence spaces $d(w, p)$ where $p \geq 1$ and $w = (w_n)_{n=1}^\infty \in c_0 \setminus \ell_1$ is a nonincreasing sequence of positive numbers with $w_1 = 1$ [6]. However, a complete classification of Banach spaces with a unique symmetric basis seems far from being achieved.

In [8], Kalton investigated Orlicz sequence spaces when the restriction of local convexity is lifted and proved that if $\lim_{t \rightarrow 0} F(t)/t = \infty$ then ℓ_F has a unique unconditional basis (hence, a unique symmetric basis). The ℓ_p spaces for $0 < p < 1$ belong to this class. Nawrocki and Ortyński [18] studied the quasi-Banach Lorentz sequence spaces $d(w, p)$ for $0 < p < 1$, described their Banach envelopes, and showed that all symmetric bases of $d(w, p)$ for $0 < p < 1$ are equivalent (see also [19]). The question of uniqueness of unconditional basis in $d(w, p)$ for $0 < p < 1$, on the other hand, was initiated in [18], continued in [12], and has been recently completely settled in [4].

All the above mentioned spaces have a 1-symmetric, and so 1-unconditional, basis which induces a p -convex lattice structure for some $p > 0$. If a quasi-Banach space X is isomorphic to a closed subspace of a p -convex quasi-Banach lattice, then X is also p -convex and it is called *natural* (see [9]).

Our aim is to prove that all natural quasi-Banach spaces with symmetric basis whose Banach envelope is isomorphic to ℓ_1 have a unique symmetric basis. This turns out to be the most important case since the Banach envelope of a natural quasi-Banach space with a basis contains ℓ_1^n 's uniformly complemented [10]. Let us point out that if a quasi-Banach space is not natural then its Banach envelope can behave very differently, as shown in [7, Sections 6 & 7].

Throughout this article we use standard Banach space theory terminology and notation, as may be found in [1,16]. For the necessary background on quasi-Banach spaces we refer the reader to the books [11,20] and the paper [13].

* Corresponding author.

E-mail address: fernando.albiac@unavarra.es (F. Albiac).

2. The main result

Suppose that $(x_n)_{n=1}^\infty$ is a symmetric basis with symmetric constant K of a quasi-Banach space $(X, \|\cdot\|)$ and assume that the Banach envelope of X , $(\hat{X}, \|\cdot\|_c)$ is isomorphic to $(\ell_1, \|\cdot\|_1)$. We observe that then $(x_n)_{n=1}^\infty$ is also a K -symmetric basis of \hat{X} and that

$$K^{-1} = K^{-1} \|x_n\| \leq \|x_n\|_c \leq \|x_n\| = 1, \quad n \in \mathbb{N}.$$

That is, $(x_n)_{n=1}^\infty$ is a semi-normalized symmetric basis of \hat{X} equivalent to the normalized basis $(x_n/\|x_n\|_c)_{n=1}^\infty$. Since ℓ_1 has unique symmetric basis, $(x_n)_{n=1}^\infty$ is equivalent to the canonical ℓ_1 -basis, i.e., there is a constant $D > 0$ so that

$$D^{-1} \sum_{n=1}^\infty |\alpha_n| \leq \left\| \sum_{n=1}^\infty \alpha_n x_n \right\|_c \leq D \sum_{n=1}^\infty |\alpha_n|,$$

for any sequence of scalars $(\alpha_n)_{n=1}^\infty \in c_{00}$.

Our main result, Theorem 2.2, relies upon the following lemma. The technique of finding “uniformly large coefficients” was introduced by Kalton in [8] to prove the uniqueness of unconditional basis in nonlocally convex Orlicz sequence spaces and was extended to the framework of quasi-Banach lattices in [12]. It has become crucial to further determine the uniqueness of unconditional basis in other quasi-Banach spaces (cf. [2,3,12,14]).

Lemma 2.1. (See [12, Theorem 2.3].) Let X be a natural quasi-Banach space with two normalized unconditional bases $(x_n)_{n=1}^\infty$ and $(y_n)_{n=1}^\infty$. Let $(x_n^*)_{n=1}^\infty$ and $(y_n^*)_{n=1}^\infty$ be the sequences of biorthogonal linear functionals associated to $(x_n)_{n=1}^\infty$ and $(y_n)_{n=1}^\infty$, respectively. Set $S \subset \mathbb{N}$ and suppose that there are a constant $\beta > 0$ and an injective map $\sigma : S \rightarrow \mathbb{N}$ so that

$$|x_{\sigma(n)}^*(y_n)| \geq \beta, \quad n \in S.$$

Then, there is a constant $\rho > 0$ such that for any scalars $(a_n)_{n=1}^\infty \in c_{00}$,

$$\rho \left\| \sum_{n \in S} a_n x_{\sigma(n)} \right\| \leq \left\| \sum_{n \in S} a_n y_n \right\|.$$

Theorem 2.2. Let X be a natural quasi-Banach space with normalized symmetric basis $(x_n)_{n=1}^\infty$ and whose Banach envelope is isomorphic to ℓ_1 . Suppose that $(y_n)_{n=1}^\infty$ is another normalized symmetric basis of X . Then there exist positive constants Δ_1, Δ_2 so that

$$\Delta_1 \left\| \sum_{n=1}^\infty a_n x_n \right\| \leq \left\| \sum_{n=1}^\infty a_n y_n \right\| \leq \Delta_2 \left\| \sum_{n=1}^\infty a_n x_n \right\|, \quad (2.1)$$

for any choice of scalars $(a_n)_{n=1}^\infty \in c_{00}$.

Proof. The bases $(x_n)_{n=1}^\infty$ and $(y_n)_{n=1}^\infty$ are symmetric bases in \hat{X} , both equivalent to the canonical basis of ℓ_1 . Let $D > 0$ such that

$$\frac{1}{D} \sum_{n=1}^\infty |\alpha_n| \leq \left\| \sum_{n=1}^\infty \alpha_n x_n \right\|_c \leq D \sum_{n=1}^\infty |\alpha_n|,$$

whenever $\sum_{n=1}^\infty \alpha_n x_n \in X$, and $C > 0$ so that

$$\frac{1}{C} \sum_{n=1}^\infty |\alpha_n| \leq \left\| \sum_{n=1}^\infty \alpha_n y_n \right\|_c \leq C \sum_{n=1}^\infty |\alpha_n|,$$

whenever $\sum_{n=1}^\infty \alpha_n y_n \in X$.

Put

$$x_k = \sum_{n=1}^\infty y_n^*(x_k) y_n, \quad k = 1, 2, \dots,$$

and

$$y_n = \sum_{k=1}^\infty x_k^*(y_n) x_k, \quad n = 1, 2, \dots$$

For each fixed $k \in \mathbb{N}$ we have

$$\begin{aligned}
1 &= |x_k^*(x_k)| \leq \sum_{n=1}^{\infty} |y_n^*(x_k)x_k^*(y_n)| \leq \sup_n |x_k^*(y_n)| \sum_{n=1}^{\infty} |y_n^*(x_k)| \leq C \sup_n |x_k^*(y_n)| \left\| \sum_{n=1}^{\infty} y_n^*(x_k)y_n \right\|_c \\
&= C \sup_n |x_k^*(y_n)| \|x_k\|_c \leq C \sup_n |x_k^*(y_n)|.
\end{aligned}$$

Thus, we can define a function $f: \mathbb{N} \rightarrow \mathbb{N}$ by $f(k) = n(k)$, where $n(k)$ is such that $|x_k^*(y_{n(k)})| > 1/2C$. Of course, f need not be one-to-one nor onto, but it is not far from being injective. Indeed, given $n_0 \in \mathbb{N}$ suppose there are integers k_1, \dots, k_N so that

$$|x_{k_1}^*(y_{n_0})| > 1/2C, \quad \dots, \quad |x_{k_N}^*(y_{n_0})| > 1/2C.$$

Then,

$$\frac{N}{2C} \leq \sum_{k=1}^N |x_{k_i}^*(y_{n_0})| \leq D \left\| \sum_{k=1}^N x_{k_i}^*(y_{n_0})x_{k_i} \right\|_c = D \|y_{n_0}\|_c \leq D.$$

Hence there is a partition of \mathbb{N} into at most $N \leq 2DC$ disjoint subsets A_1, \dots, A_N so that the restriction of f to each A_i is one-to-one. We pick one of these sets of the same cardinality as \mathbb{N} , say A_1 , and let $S = f(A_1)$. This way, the map $f: A_1 \rightarrow S$ is a bijection. Let us consider its inverse,

$$\sigma: S \rightarrow A_1, \quad n \rightarrow \sigma(n) \quad \text{so that} \quad |x_{\sigma(n)}^*(y_n)| > 1/2C.$$

By appealing to Lemma 2.1 we find $\rho > 0$ so that for any $(a_n)_{n=1}^{\infty} \in c_{00}$ we have

$$\rho \left\| \sum_{n \in S} a_n x_{\sigma(n)} \right\| \leq \left\| \sum_{n \in S} a_n y_n \right\|. \quad (2.2)$$

The symmetry of the bases yields positive constants Γ_1, Γ_2 such that

$$\frac{1}{\Gamma_1} \left\| \sum_{n=1}^{\infty} a_n x_n \right\| \leq \left\| \sum_{n \in S} a_n x_{\sigma(n)} \right\| \leq \Gamma_1 \left\| \sum_{n=1}^{\infty} a_n x_n \right\| \quad (2.3)$$

and

$$\frac{1}{\Gamma_2} \left\| \sum_{n=1}^{\infty} a_n y_n \right\| \leq \left\| \sum_{n \in S} a_n y_n \right\| \leq \Gamma_2 \left\| \sum_{n=1}^{\infty} a_n y_n \right\|, \quad (2.4)$$

for any $(a_n)_{n=1}^{\infty} \in c_{00}$. Combining (2.2)–(2.4) we obtain the left-hand side inequality of Theorem 2.2 with $\Delta_1 = (\Gamma_1 \Gamma_2)^{-1} \rho$. The right-hand side inequality in (2.1) follows by interchanging the roles of the two bases by a symmetric argument, and this completes the proof. \square

Concluding remarks. (a) The hypotheses of Theorem 2.2 are not sufficient to ensure the uniqueness of *unconditional* basis in X . As a matter of fact, Kalton [8] found examples of nonlocally convex Orlicz sequence spaces whose Banach envelope is ℓ_1 and, however, do not have a unique unconditional basis.

(b) There are natural quasi-Banach spaces whose Banach envelope is not ℓ_1 with unique symmetric basis. Take, for instance, $X = d(w, p)$ for $0 < p < 1$, where the sequence of weights $(w_n)_{n=1}^{\infty}$ satisfies the condition

$$\inf_n \frac{(w_1 + \dots + w_n)^{1/p}}{n} = 0.$$

Then, by [18, Proposition 3], X has unique symmetric basis, but by [18, Theorem 1], the Banach envelope of X is not isomorphic to ℓ_1 .

(c) A normalized basis is *perfectly homogeneous* if it is equivalent to all its normalized block basic sequences, that is, perfectly homogeneous bases are a special case of symmetric bases. For Banach spaces, a classical result of Zippin [21] proved that perfectly homogeneous bases are equivalent to either the canonical c_0 -basis or the canonical ℓ_p -basis for some $1 \leq p < \infty$. In [5] the authors show that Zippin's techniques can be extended to prove that perfectly homogeneous bases of nonlocally convex quasi-Banach spaces are always equivalent to the canonical ℓ_p -basis for some $0 < p < 1$.

Our work leaves open the following question.

Problem 2.3. Are there any nonlocally convex quasi-Banach spaces with symmetric basis without the property of uniqueness of symmetric basis?

Acknowledgments

The authors would like to acknowledge support from the Spanish Ministerio de Educación y Ciencia Research Project *Espacios Topológicos Ordenados: Resultados Analíticos y Aplicaciones Multidisciplinares*, reference number MTM2007-62499.

References

- [1] F. Albiac, N.J. Kalton, Topics in Banach Space Theory, Grad. Texts in Math., vol. 233, Springer-Verlag, New York, 2006.
- [2] F. Albiac, N. Kalton, C. Leránoz, Uniqueness of the unconditional basis of $l_1(l_p)$ and $l_p(l_1)$, $0 < p < 1$, Positivity 8 (4) (2004) 443–454.
- [3] F. Albiac, C. Leránoz, Uniqueness of unconditional basis of $l_p(c_0)$ and $l_p(l_2)$, $0 < p < 1$, Studia Math. 150 (1) (2002) 35–52.
- [4] F. Albiac, C. Leránoz, Uniqueness of unconditional basis in Lorentz sequence spaces, Proc. Amer. Math. Soc. 136 (5) (2008) 1643–1647.
- [5] F. Albiac, C. Leránoz, On perfectly homogeneous basis in quasi-Banach spaces, submitted for publication.
- [6] Z. Altshuler, P.G. Casazza, Bor Luh Lin, On symmetric basic sequences in Lorentz sequence spaces, Israel J. Math. 15 (1973) 140–155.
- [7] F. Cabello Sánchez, J.M.F. Castillo, P.L. Papini, Seven views on approximate convexity and the geometry of K -spaces, J. London Math. Soc. 72 (2005) 457–477.
- [8] N.J. Kalton, Orlicz sequence spaces without local convexity, Math. Proc. Cambridge Philos. Soc. 81 (2) (1977) 253–277.
- [9] N.J. Kalton, Convexity conditions on non-locally convex lattices, Glasg. Math. J. 25 (1984) 141–152.
- [10] N.J. Kalton, Banach envelopes of non-locally convex spaces, Canad. J. Math. 38 (1986) 65–86.
- [11] N.J. Kalton, N.T. Peck, J.W. Rogers, An F -Space Sampler, London Math. Soc. Lecture Notes, vol. 89, Cambridge Univ. Press, Cambridge, 1985.
- [12] N.J. Kalton, C. Leránoz, P. Wojtaszczyk, Uniqueness of unconditional bases in quasi-Banach spaces with applications to Hardy spaces, Israel J. Math. 72 (3) (1990) 299–311.
- [13] N.J. Kalton, Quasi-Banach spaces, in: Handbook of the Geometry of Banach Spaces, vol. 2, North-Holland, Amsterdam, 2003, pp. 1099–1130.
- [14] C. Leránoz, Uniqueness of unconditional bases of $c_0(l_p)$, $0 < p < 1$, Studia Math. 102 (3) (1992) 193–207.
- [15] J. Lindenstrauss, L. Tzafriri, On Orlicz sequence spaces, Israel J. Math. 10 (1971) 379–390.
- [16] J. Lindenstrauss, L. Tzafriri, Classical Banach Spaces. I. Sequence Spaces, Springer-Verlag, Berlin, 1977.
- [17] J. Lindenstrauss, M. Zippin, Banach spaces with a unique unconditional basis, J. Funct. Anal. 3 (1969) 115–125.
- [18] M. Nawrocki, A. Ortyński, The Mackey topology and complemented subspaces of Lorentz sequence spaces $d(w, p)$ for $0 < p < 1$, Trans. Amer. Math. Soc. 287 (2) (1985) 713–722.
- [19] N. Popa, Basic sequences and subspaces in Lorentz sequence spaces without local convexity, Trans. Amer. Math. Soc. 263 (2) (1981) 431–456.
- [20] S. Rolewicz, Metric Linear Spaces, second ed., Math. Appl. (East European Series), vol. 20, D. Reidel Publishing Co., Dordrecht, 1985.
- [21] M. Zippin, On perfectly homogeneous bases in Banach spaces, Israel J. Math. 4 (1966) 265–272.